

POST-ABORTION SURVIVOR SYNDROME: SIGNS AND SYMPTOMS

Philip G. Ney, MD, FRCP(C), MA, Claudia K Sheils, RN
Marek Gajowy, MA

ABSTRACT: Clinical observations indicated that those psychiatric patients who survived when a preborn sibling died were adversely affected by being allowed to live. It seemed being a survivor of a pregnancy loss, particularly abortion, contributed to psychiatric illnesses. Data was collected from a sample of 293 adults - 98 patients and 195 counseling trainees. A self-report questionnaire with visual analogue, rating, and descriptive questions was used to ascertain the extent of common psychiatric symptoms. These were analyzed to determine if there were any significant associations with various types of pregnancy outcome. Correlations and stepwise regression analyses demonstrated a cluster of existential symptoms for those surviving when their preborn siblings were aborted. The symptom that most closely associated with the “total number of abortions in my mother’s first pregnancy” was, “I feel I don’t deserve to be alive.” There were different and more loosely clustered symptoms found in patients whose mother miscarried.

Conclusion: there is a reasonably definable syndrome of symptoms in patients associated with the abortion of their sibling, which we have termed, the Post Abortion Survivor Syndrome.

KEYWORDS: Post-Abortion Survivor Syndrome, pregnancy loss, abortion, siblings, suicide, existential guilt, impending doom.

INTRODUCTION

There appears to be a paradoxical response when, for reasons over which they have no control, a person’s life is spared when those who are near and are dear to them die. You might think that people who survive should be glad to be alive and greet every dawn with gladness. Instead, studies have shown many survivors of torture, concentration camps, disasters, accidents, and illnesses have a pervasive sense of guilt, morbid thoughts, suicidal ideation, and difficulty grappling with the exigencies of life. What was first known as the “concentration camp syndrome”(Chodoff, 1963) later became the “survivor syndrome (Niederland, 1968).” Symptoms included cognitive and memory disturbances, depression and survivor guilt, chronic anxiety related to the fear of renewed persecution, and phobic fears. Frequently there are sleep disturbances including insomnia, nightmares, and anxiety dreams related to persecution, as well as somatic manifestations. Dr. Wanda Poltawska (1989) wrote *And I am Afraid of my Dreams* in which she describes terrifying night visions which persisted for many years until she was able to record and talk about her gruesome experience as a “guinea pig” in so-called medical experiments by the Nazis in Ravensbrück.

Work done while with the Department of Family Practice, Faculty of Medicine, University of British Columbia. Contact: Dr. Philip G. Ney, Box 27103, Victoria, B.C. Canada, V9B 5S4
Phone: (250) 642 1848 Fax: 250 642 1841 Email: pgney@telus.net Institutional Review Board Approval: University of British Columbia, Research Ethics Committee
No experimental intervention.

Holocaust survivors have indicated loss of childhood memories, distorted perceptions of personal identity, and difficulties with interpersonal relationships (Kaminer & Lavie, 1994). It soon became apparent that these symptoms not only affected the survivors, but also their children (Barocos & Barocos, 1973; Danieli, 1981; Krell, 1979; Kestenberg, 1990; Roland-Klien & Dunlop, 1997). The survivors' children had symptoms resembling those of their parents, including depression, anxiety, phobias, guilt and separation problems, frightening dreams, and environmental misperceptions. The children's depression was attributed to anger turned inwards. Those children were raised to view the world as a dangerous place, but no expression of aggression was tolerated at home.

According to Krell (1979) the terrifying experiences are inevitably transmitted to offspring whether these are discussed openly or as veiled references or mysterious outbursts of grief. The parent's lack of communication might contribute to the child's increased depression because of his/her preoccupation with fantasies about what his/her parent may have experienced. These fantasies may be more frightening and pathogenic than the actual events. Miller (1995) believes that children unconsciously re-enact their parents' fate all the more intensely the less precise their knowledge of it. Wardi (1992) believes children of Holocaust survivors may be discharging their parents' unresolved, unconscious conflicts. I have hypothesized that the trans-generational re-enactment of unresolved conflicts arising from trauma is an adaptive mechanism whereby the mind is forcing the individual to confront the subconscious problem so that, once resolved, the brain can function more efficiently and allow the individual to be successful in communication and work (Ney, 1988, 1989).

Holocaust survivors were found to be vulnerable to a reactivation of this "survivor syndrome" when there was subsequent trauma, even though they appeared to be asymptomatic at the time (Robinson, Hemmendinger, Netanel, et al, 1994). Vulnerability to psychological distress also affected second-generation Holocaust survivors when faced with trauma such as breast cancer (Baider, Peretz, Hadani, et al, 1994). It was found that, although there were some differences, most of these symptoms could be found in the survivors of torture (Wenzel, Griengl, Stompe, Mirzaei, & Kiefer, 2000), those living in a war torn country (Bramsen & van der Ploeg, 1999), those surviving planned explosions (North, Nikon, Shariat, et al, 1999), and those surviving unplanned sinking (Bolton, O'Ryan, Udwin, Boyle, & York, 2000).

Holocaust survivors have indicated loss of childhood memories, distorted perceptions of personal identity, and difficulties with interpersonal relationships (Kaminer & Lavie, 1994). It soon became apparent that these symptoms not only affected the survivors, but also their children (Barocos & Barocos, 1973; Danieli, 1981; Krell, 1979; Kestenberg, 1990; Roland-Klien & Dunlop, 1997). The survivors' children had symptoms resembling those of their parents, including depression, anxiety, phobias, guilt and separation problems, frightening dreams, and environmental misperceptions. The children's depression was attributed to anger turned inwards. Those children were raised to view the world as a dangerous place, but no expression of aggression was tolerated at home.

According to Krell (1979) the terrifying experiences are inevitably transmitted to offspring whether these are discussed openly or as veiled references or mysterious outbursts of grief. The parent's lack of communication might contribute to the child's increased depression because of his/her preoccupation with fantasies about what his/her parent may have experienced. These fantasies may be more frightening and pathogenic than the actual events. Miller (1995) believes that children unconsciously re-enact their parents' fate all the more intensely the less precise their knowledge of it. Wardi (1992) believes children of Holocaust survivors may be

discharging their parents' unresolved, unconscious conflicts. I (PGN) have hypothesized that the trans-generational re-enactment of unresolved conflicts arising from trauma is an adaptive mechanism whereby the mind is forcing the individual to confront the subconscious problem so that, once resolved, the brain can function more efficiently and allow the individual to be successful in communication and work (Ney, 1988, 1989).

Holocaust survivors were found to be vulnerable to a reactivation of this "survivor syndrome" when there was subsequent trauma, even though they appeared to be asymptomatic at the time (Robinson, Hemmendinger, Netanel, et al, 1994). Vulnerability to psychological distress also affected second-generation Holocaust survivors when faced with trauma such as breast cancer (Baider, Peretz, Hadani, et al, 1994). It was found that, although there were some differences, most of these symptoms could be found in the survivors of torture (Wenzel, Griengl, Stompe, Mirzaei, & Kiefer, 2000), those living in a war torn country (Bramsen & van der Ploeg, 1999), those surviving planned explosions (North, Nikon, Shariat, et al, 1999), and those surviving unplanned sinking (Bolton, O'Ryan, Udwin, Boyle, & York, 2000).

Clinical impressions of Holocaust survivors seem to indicate that, although there were symptoms of high anxiety and deep depression, there were more existential type symptoms that could not easily be used in determining an ICD or DSM diagnosis. These symptoms appeared to arise from conflicts that the person had been struggling with for long periods. Although patients referred to me for consultation had received various combinations of medication and psychotherapy, most felt that their deeper dilemmas had not been understood. Although the term survivor has frequently been used in a very broad context to include anybody that might have died for any reason, it is used in this paper to indicate those who remain alive after some force over which they had no control prematurely ended the life of somebody near and dear to them, namely a preborn sibling.

Approximately 30 years ago, an eight-year-old girl, referred to me for evaluation, forced me to consider the impact of being a pregnancy loss survivor. The mother who brought this, her only child, was worried because she was not sleeping well, was irritable, could not concentrate, frequently burst into tears, and often seemed to be preoccupied. This all began with a recurrent nightmare. The child clearly described her very frightening dream to me. With three siblings, she had gone to play in a bank of sand. They tunneled into the sand and her three siblings had crawled in. The sand had collapsed and buried them alive. She could tell me very little about these children, except that she was absolutely convinced that they were her brother and sisters. Later, the mother told me she had three early miscarriages that her daughter could never have known about. It seemed to me that somehow the little girl knew or suspected the deaths of her siblings and was now worried something might happen to her. When I voiced these thoughts, the child indicated she felt she was being understood. With her mother's reassurance and an unconditional commitment to love and protect her, the child's fears and her symptoms rapidly subsided.

Having been alerted to the effect in children of surviving a pregnancy loss, I found others who knew or guessed they had lost a sibling by stillbirth, miscarriage, or abortion. When questioned their parents it appeared that these children had a surprisingly accurate knowledge their mother's pregnancy outcome.

Since the description of the Post-Traumatic Stress Disorder, many authors have attempted to determine whether survivor symptoms fit that particular diagnosis. There seemed to be less interest in the symptoms that were found early in the studies of survivors, namely existential symptoms, survivor guilt, not wishing to be alive, etc. North et al., in their study of the Oklahoma City bombing survivors,² found 45% exhibited symptoms of the Post-Disaster Psychiatric Disorder (PDPD) and 34.3% of PTSD. They suggested that the best way to screen for those who would later show evidence of psychiatric distress was by identifying the symptoms of avoidance and numbing. Intrusive re-experience and hyper arousal were symptoms so universal they were not seen to be associated with psychopathology or impairment of function.

Clinical impressions of Holocaust survivors seem to indicate that, although there were symptoms of high anxiety and deep depression, there were more existential type symptoms that could not easily be used in determining an ICD10 or DSM IVR diagnosis. These symptoms appeared to arise from conflicts that the person had been struggling with for long periods. Although patients referred to me for consultation had received various combinations of medication and psychotherapy, most felt that their deeper dilemmas had not been understood. The question herein addressed is , are there characteristic symptoms and is there a constellation of signs and symptoms sufficiently specific to constitute a syndrome for those who survived abortion?

Although the term survivor has frequently been used in a very broad context to include anybody that might have died for any reason, it is used in this paper to indicate those who remain alive after some force over which they had no control prematurely ended the life of somebody near and dear to them, namely a preborn sibling.

Approximately 30 years ago, an eight year old girl, referred to me for evaluation forced me to consider the impact of being a pregnancy loss survivor. The mother who brought this, her only child, was worried because she was not sleeping well, was irritable, could not concentrate, frequently burst into tears, and often seemed to be preoccupied. This all began with a recurrent nightmare. The child clearly described her very frightening dream to me. With three siblings, she had gone to play in a bank of sand. They tunnelled into the sand and her three siblings had crawled in. The sand had collapsed and buried them alive. She could tell me very little about these children, except that she was absolutely convinced that they were her brothers and sisters. Later, the mother told me she had three early miscarriages that her daughter could never have known about. It seemed that somehow the little girl knew or suspected the deaths of her siblings and was now worried something might happen to her. When I voiced these thoughts, the child indicated she felt she was being understood. With her mother's reassurance and an unconditional commitment to love and protect her, the child's fears and her symptoms rapidly subsided.

Having been alerted to the effects in children surviving a pregnancy loss, I found others who knew or guessed they had lost a sibling by stillbirth, miscarriage, or abortion. When I questioned their parents it appeared that these children had a surprisingly accurate knowledge of their mother's pregnancy outcome.

METHOD

The present study is an extension of a research project whose protocol and procedures were approved by the UBC medical ethics.³ The sample was composed of 293 adults, 85% women

and 13% men. Of these, 56.4 % were married, 23.4 % single, 11.1% divorced. The rest were separated, widowed, remarried, or in some other type of relationship. The age range was 22 to 75 years, mean 40.63 years. The average number of children is 2.07. The average number of hospitalisations which include giving birth is 3.61. According to Statistics Canada, this sample is reasonably representative of the population. Ninety-eight in the sample were outpatients without major psychiatric aberrations. The other 195 were relatively healthy adults training to be group counsellors.

An 86-item self-report questionnaire was filled out by the subjects, following a brief explanation by the research assistant. There were 10 questions regarding demographics, 12 short answer descriptive questions, 2 blank tables regarding all the pregnancy outcomes of the subject's mother's and the subject, 10 questions that required check marks, and 51 questions with visual analogue scales. The visual analogue scale is an 8 cm. line across which the subject is asked to draw a mark indicating his/her estimation of their position between two extremes, e.g. 'always' and 'never'. The responses were coded as a 8 point scale with additional codes for not applicable and no response. The internal reliability of the questionnaire was determined by parallel form questions. Many visual analogue questions were the same as those used in our study of the effects of pregnancy outcome on health, where the reliability and validity were determined to be adequate. (Ney PG, Wickett AR, Fung T, 1992, Ney PG, Fung T, Wickett AR 1993, ²¹

The hypotheses we tested were: 1) There is a characteristic constellation of signs symptoms sufficiently specific to constitute a syndrome for those who survived an abortion. People whose mother have a pregnancy loss are significantly affected by it. 2) a) The symptoms of those who sibling died by abortion are significantly different from those survive a siblings miscarriage. b) The existential dilemmas that arise from being an abortion survivor create unique symptoms. .

RESULTS

The questionnaire made it possible to analyze the data as one of three variables for each pregnancy outcome. For example for miscarriages: a) the total number of miscarriages of the mother of the subject's first or second, third etc. pregnancy, b) the total number of miscarriages for the subject's mother for any pregnancy up to 9, c) the total number of people whose mother had at least one miscarriage. These correlated with significance $p < 000$ in all instances. Because of these highly significant correlations, it was possible to use any of them as dependent variables for the same analysis.

The frequency of full term normal birth weight pregnancies and pregnancies that end in abortion in our sample, as indicated approximate those of the general Canadian population.

To determine the validity of the subject's perception of outcome of their mother's pregnancies we asked them how they found out (Table 1). Most indicated they were told what happened by their mothers. We also asked the subjects how their children found out about the outcome of their own pregnancies. There is spread similar to the subject's mother, with the most frequent response being, "I (subject) told them" (Table 2).

To help establish the validity of our questionnaire we analysed the symptoms that one would expect clinically to be associated with the presence of suicidal thoughts and repeated depressions. The significant associations are appropriate and are mainly those of the hypothesized post abortion syndrome. (Table 3)

A stepwise regression analysis of the visual analogue scores to the 16 questions regarding common psychiatric symptoms indicated that the closest association to the dependent variable “The total number of abortions in the first pregnancy of my mother” was “I feel I don’t deserve to be alive” (Unstandardized Coefficient B 1.171, $t = 2.047$, $p = 0.042$). If the dependent variable was the total number of subjects whose mother had at least one abortion, the most prominent symptom was “I feel that life is not worth living” ($t = -2.177$, Sig. $p < .030$).

Using the symptom “I feel I don’t deserve to be alive” as a dependent variable, a stepwise regression analysis showed that the other symptoms most closely associated to it were those arising from conflicts regarding life, death and violence to one’s self (Table 4). This sense of not deserving life appears not to be associated with any other pregnancy outcome of the subjects’ mothers’ first pregnancy (Table 5). There are predominately negative associations to existential conflicts with other types of pregnancy losses.

When the 16 symptoms are cross-correlated, controlling for age, sex, marital status, and number of children, the symptom “I don’t feel I deserve to be alive” significantly correlates with the other symptoms hypothesised as being part of the Post Abortion Survivor Syndrome (Table 6). This analysis shows those symptoms which most closely correlate with existential guilt (“I feel I don’t deserve to be alive”) are similar to the results of the stepwise regression analysis, namely “I am not glad to be alive”, “I am not pleased with who I am”, “I sense something terrible is going to happen to me”, and “I have thoughts I can’t control”. The symptom that most closely correlates with “I feel life is worth living” is “I feel sad”. The other symptoms correlate with each other, as one would expect clinically. “I hear or see or feel things that appear from nowhere” (hallucinations) correlates most highly with “I have feelings that things are unreal” (derealisation), and “I fear I am losing my mind” (pre-psychotic panic).

When questions about the mother’s abortion or the chance of being aborted are asked separately there are somewhat different closely associated symptoms but all are part of what could be called the Post Abortion Survivor Syndrome, (Table 7).

We did an analysis to determine the most prominent symptoms of those who survived when unborn siblings were miscarried. We found that the only two symptoms of significance, which ever why we asked the question, statistically were: “I feel life is worth living” ($p < 0.000$ to 0.037) and “I feel angry” ($p < 0.001$). There were no significant associations of any of the 16 symptoms for premature births, still births or early infant deaths.

If the dependent variable is the subject’s repeated depressions, the most closely associated possible contributors were the impact of mistreatment on the early life of the subject and the total number of persons who had at least one abortion. When the dependent variable is “The total number of abortions in my or my partner’s first pregnancy” the most prominent factors are emotional or intellectual neglect in childhood, and the total number of persons whose mother had at least one abortion.

Why women choose to terminate a pregnancy is a complex and hotly debated subject. The amount they want to have a child is a notoriously poor indicator because wantedness fluctuates widely from day to day depending on mood, turmoil in a relationship, finances etc. Using a matched samples t test on those subjects whose mother had a full term pregnancy or miscarriage or abortion in her first pregnancy and those whose mothers had no full term pregnancy,

miscarriage or abortion. Assuming equal variance (confirmed by Levene's Test for Equality of Variance) we found no significant differences in the background and demographic variables. In these three groups in their age, sex, marital status, number of hospitalisations and highest level of education there was no significant difference. To determine whether or not there was a difference in the early experience of subjects in these three groups, we included "my parents were happily married", "we had a happy family life" and "my brothers and sisters are doing well" as variables. There was no significant t test difference. The only significant difference in that analysis was the symptom "I feel I don't deserve to be alive". (Std Error Difference .572, Sig. (2 tailed) .042, 95% Confidence Interval -2.297 to .045. If a separate t test is done on the two groups, a) those whose mother aborted her first pregnancy, b) those who did not, the significant differences in the symptoms were "I feel I don't deserve to be alive" and "I have injured myself"

We determined what were the problems in the abortion survivor's present life when there had been a statistically high chance of being aborted. We found there were statistically significant associations with an inability to trust others and troubling dreams. When the dependent variable was the total number of abortions of the subject's mothers, we found low self esteem and feelings of being haunted were significantly associated.

Table 8 shows the past problems the abortions survivors have had. There were no significant problems in the subject's past associated with miscarriages in the subject's mother's first pregnancy. There were no significantly associated past problems for those whose mother never had an abortion but had some other pregnancy loss.

To determine whether there was a trans-generational component to abortion, we calculated the number of abortions for the subject's mother in three ways: 1) the total number of abortions in the mothers' first pregnancy, 2) the total number of abortions in any pregnancy for all the mothers, 3) the total number of women who had one or more abortions. These three correlate closely. They tend to indicate the subject had a reasonably accurate awareness of their mother's pregnancy outcomes (Table 9). There is a significant correlation between the subject or the subject's partner aborting her first pregnancy and his/her mother aborting her first pregnancy. There are significant correlations between a subject or a subject's partner aborting her first and second pregnancies and his/her mother aborting her first and second pregnancies.

Although subjects are affected by other kinds of pregnancy loss, they currently feel more affected by their mother's abortion. Though there is greater variability, the unstandardized coefficient indicates that subjects are next more deeply affected by their mother's stillbirth. Other regression analyses show that subjects were less inclined to talk to their mother about her abortion than about other losses. They also indicated they were less likely to have completed grieving the loss of a sibling through abortion.

It appears that for men, the symptoms associated with their mother's abortions are different from those of the whole group. These symptoms appear to be more frequently violent, self-destructive tendencies and fears of impending doom.

Factors, as they are calculated by standard statistical techniques, derived from visual analogue scales are represented here as a combination of the frequency and the severity of the impact on

the individual. When this is done the most prominent symptoms of PASS are of an existential type: (Table 10)

DISCUSSION

From these results it appears that those who are affected by the loss of their unborn siblings to pregnancy termination have predominantly existential type symptoms. The sense that they do not deserve to be alive correlates closely with other symptoms of life and death conflicts: a sense of impending doom, self-destructiveness and a high level of anxiety that would make them feel that they are losing their mind. Although some people appeared to cope with their existential conflicts, a significant number of others indicated repeated depression and psychiatric hospitalisation in the past. One would have to wonder whether the need for repeated psychiatric treatment would have occurred had the existential conflicts been addressed earlier.

There are symptoms associated with the subject's mother having had one or more miscarriages.. However there is a marked difference in the symptoms of abortion survivors. These seem to be more likely to arise from existential conflicts regarding life, living and the meaning of existence which are not detected by clinicians who leave questions regarding them from their usual patient evaluation.

The question whether the abortion survivors come from a different demographic and family background seems to have been answered by the data for those whose mother's had full term babies or miscarriages or abortions. According to the independent samples T there appears to be no detectable difference. This suggests that it was not their genetic predisposition or quality of family life but that their mother had an abortion rather than a full term baby or a miscarriage. The next study will include more information from the mother and father of the subjects. When the dependent variable is a variety of other questions regarding the subject's fear that he/she might have also been terminated or that his/her parents considered aborting him/her, there are feelings of unreality and a tendency to suicide. Whether the analysis is a stepwise regression analysis or a partial correlation, there is reasonable uniformity in the constellation of symptoms.

As a comparison, symptoms of those who lost siblings through miscarriage are quite different. Interestingly, although they feel angry, they feel that life is worth living and they have no trouble with interpersonal relationships. One of the more prominent symptoms of the abortion survivors is that they are not glad to be alive whereas those who survived when their mothers had miscarriages indicate they are glad to be alive. Why abortion survivors should have more difficult symptoms than miscarriage survivors may be related to the fact that their unborn siblings died quite differently. Abortion survivors have more reason to believe they may have contributed to the death of a sibling since this death was determined by a conscious choice of their parents. This is especially the case if the survivor was a "difficult child", for then the parents may have decided they don't want any more children lest they have another "like this one". Those who survive where there has been a miscarriage of a sibling survive an unplanned incident. They can, without self-recrimination, be glad that they are alive. .

In addition to determining symptoms, we asked the subjects to respond to questions regarding their past and present interpersonal and psychological problems. Depending on how the question regarding subjects' mothers' abortion is asked, the current problems varied somewhat. Those subjects who are abortion survivors indicate their mother's abortion lowers

their self-esteem and inclines them to feel that they are being haunted; conceivably this sensation maybe an image or a fantasy of the aborted child.

We found there was a tendency on the part of women whose mother had an abortion to have an abortion themselves. The total number of abortions for the subject or the subject's partner best correlates with the neglect during childhood and the abortions of their mothers. Kent, Greenwood and Nicholls²⁴ found that women who aborted were often carrying out their parents' unconscious wish to abort them.

Abortion survivors not only survive the loss of one or more of their siblings, but may also live through the trauma of their parent's distress following an abortion and possibly childhood mistreatment. One patient described this very succinctly, "How could you, (parents), be loving to me and yet have killed one of my siblings. You still might do something to me. I do not trust you. I do not trust the anger I feel towards you. I sometimes want to kill you. Yet I need you. It is safer if I can see and observe you all the time. I will do that until I am old enough to run away."⁴(Ney, PG,,Peeters MA 1996)

Subjects who had past problems with repeated depression indicate that possible causes were mistreatment as a child and the abortions of their mothers. The mother's abortion was also a prominent possible contributor to the subject or the subject's partner having an abortion.

It appears that the symptoms and problems of abortion survivors have a number of distinctions from those of other types of survivors. This is not surprising when it is understood that the children feel threatened by those that are supposed to care for them in any situation. Most children have probably heard many stories of parents sacrificing themselves for their children. When children realize that their parents were prepared to sacrifice one of their own children, it is understandable these subjects may have deep fears of those who are close to them and wish to care for them.

There are bound to be some similar symptoms in PTSD and PASS because the etiologies have common features. The DSM IV TR notes that PTSD symptoms arise following exposure to an "extreme stressor" including "witnessing a dead body or body parts". This could occur in the life of an abortion survivor on seeing for the first time a picture of an aborted fetus and realizing her/his sibling could have looked like that. However the same could be true of a person whose twin was miscarried. The evidence presented here seems to indicate that miscarried survivors are not so adversely effected by such an event that they develop symptoms anything like those of an abortion survivor. In clinical practice, miscarriage survivors on learning of a sibling lost in utero, tend to show interest rather than horror or disgust. When they see a picture of an aborted fetus, they express distress but this tends no to last like the symptoms of PTSD.

The discovery of being an abortion survivor, seems to be a gradual dawning of awareness rather than the sudden extreme stressor in PTSD. Both PASS and PTSD patients have a sense of "impending doom" or "foreshortening future". They both have nightmares but PTSD sufferers tend to visually re-experience the trauma while PASS people tend to see very abstract scenes which symbolically represent their intense conflicts. Both groups have guilt on surviving when others did not, but post abortion survivors have usually grown up with this sense. It almost feels familiar if not natural to them. Whereas PTSD are generally symptom free before being traumatized, PASS sufferers have symptoms starting in their childhood.

In both groups there is a general tendency to avoid any encounter with the traumatic stressors. For PASS, this is often their parents. Yet abortion survivors have such a fear of being abandoned when and if they become unwanted, they also show great care for parents, especially elderly mothers. At the same time they feel a deep resentment they can hardly hide. PASS

people often think of death and usually dance with death in some form but they are not genuinely suicidal like PTSD. Both groups find discussing the basic issues very painful, but PTSD sufferers seem to understand it is necessary while PASS patients find their key conflicts so complex, they give up easily.

Both groups tend to be easily irritated and unnecessarily angry. Abortion survivors may become enraged at any one who in casual conversation or in therapy broaches the subject of their mother's abortion. PTSD seem to be most angry at whoever should have protected them from the traumas but for weak reasons did not.

It is estimated that about 15% of multiples grow up as singleton survivors. There is extensive literature on the outcome for the surviving twin when there has been twin-twin transfusion syndrome in terms of neurological difficulties (Sago, Hayashi, Salto, et al, 2010)(23) but little on the psychological impact. "Twin concordance for suicide attempts more likely reflects a genetic predisposition than a behavioural reaction to the loss." (Segal, 2009)(24) This statement is being questioned in light of more recent studies. The impact of the "vanishing twin" has been considered in some scientific writings (Rinborg, Lidgaard, Freiesleben et al, 2007)(25) but this subject is much more extensively dealt with in web sites and blogs. Hayton makes a short comment on her blog "Womb Twin Survivors" (26) about abortion survivors, "I'm pretty sure my mother aborted my twin"

It is not possible to make direct comparisons to similar studies because there appear to be none. However, we can compare these results with our own data; the symptoms of those who might be affected by their mothers' miscarriage, those who were mistreated as children, and those who had an abortion or those whose partner had abortion. There is obviously an overlap of symptoms with those who are suffering from child abuse and neglect and those who are suffering the symptoms of having had an abortion, for both groups felt very exposed when they were most vulnerable.

It appears that our questions which allowed subjects to mark their past or present situation or reaction on a visual analogue scale, is a useful way of collecting data. There were few who did not understand the nature of the question and appreciate not having to categorize themselves. The analogue data, converted to an 8-point scale, lends itself well to statistical analysis.

Using data from the range of marital status in our sample compared to STATISTICS CANADA, these subjects can be considered to represent the general population. This was an unselected sample of patients and the findings can be generalized to a psychiatric practice. There were asymptomatic abortion survivors who were not experiencing significant de-compensation but in the future they might under added stress. The validity of the symptoms seems to be established by the higher correlation with the history of high psychiatric disorder, repeated depression etc.

The overlap in symptoms between the different experiences of loss or mistreatment probably indicates that in many instances there were common contributing factors. Other research in progress seems to indicate the lack of partner support is both cause and effect of a woman's abortion. In the near future we hope to determine whether there is a difference in the symptom constellation of those who were born before as opposed to those who were born after their mother's abortion.

The constellation of symptoms arising from the knowledge or strong suspicion of their mother's aborting a child, especially it seems the first pregnancy seems logical in the light of the very

difficult conflicts that arise. The abortion survivor feels guilty for existing and therefore can't enjoy life or fully use their opportunities and abilities. They may not have been told by their mother about her abortion but there is much circumstantial support to their suspicion and intuitive awareness. To be more certain, they may want to ask their mother but at the same time don't want to know for it is a most discomfoting revelation. Most survivors seem to "keep a ear open" even when their mind rejects the idea. When they do find out for sure, they are both relieved and angry; relieved because it explains so well their inner turmoil and conflicted experience and angry ("mother how could you do such a thing") because they have been deprived of a sibling with whom they could have shared their joys and sorrows.

Their existential conflicts are so intense they begin to fear they are losing their mind, particularly because there doesn't seem to be any reason they should feel so torn. They may feel guilty for not have done something to rescue their sibling or by their existing or obnoxious behaviour for creating a feeling in their parents for not wanting another child. Although they are afraid to die, they flirt with death, doing extreme sports or overdosing with drugs or alcohol. To reassure themselves they are alive, they may wish to see and taste their blood flowing from a self inflicted wound. They may sense they have lost a sibling and don't know how it may have happened. It is not surprising they have a sense of impending doom. "Whatever happened to my brother could also happen to me" Unlike the miscarriage survivors they are not glad to be alive, mainly it seems because life is so confused and painful. Over time we have developed a treatment program (Ney PG Ball K Shiels C 2010....) that seems to help abortion survivors but even with intensive psychotherapy resulting in symptomatic improvement, their key conflicts continue.

Miscarriage survivors feel glad to live and do not have the existential dilemmas of abortion survivors, yet there are some symptom similarities. The data for this may be explained by the sample having 98 outpatients. Why they are angry, we can't explain except for the possibility of their having lost a sibling they could have loved.

The transgenerational tendency of mother's to have an abortion may arise from a desire to understand why the subject's mother would choose to terminate her child by re-enacting that conflict.(Ney PG 1989) There is also evidence that unresolved conflicts of surviving parents are felt by their children, (Krell) We need to redo this type of study controlling for the impact of abortion on the subject's ability to bond and nurture her babies following an abortion. In previous research we found that there was a higher incidence or post-partum depression in women who previously had an abortion. It appeared that the depression interfered with their ability to bond to her newborn infant. The lack of a strong, resilient bond may contribute to the subsequent higher incidence of a abuse and neglect. (Kersting A Ohrmann P, 2009)

Whichever way we analysed the data, the finding that miscarriage survivors feel glad to be alive while abortion survivors feel they don't deserve to be alive, stands out. This is not easy to explain but this inadequate analogy might help. A little family of good parents and two children were taking a seaside holiday. One day while playing along the top of an unguarded cliff, one of the children tripped and fell to her death on the rocks far below. The parents berated themselves for their carelessness. However after a period of grieving, the surviving sibling rebounded to health and vigor. She was heard to remark in an unguarded moment, "I'm very sad she is gone but now I can play with her toys." Another family not far from them were walking along a path at the top of the cliff. The mother and father after a quiet discussion about never wanting two children, pushed one child to his death. The remaining sibling did not see them do it but heard

the child the child cry out as he fell. This survivor was morose for a very long time and grew up seldom enjoying life.

This study collected a great deal of information that still needs to be analysed. Like our study on the effects of pregnancy losses on a woman's health,(Ney, 2010) we need to collect data from a sample of patients, men and women, who are waiting to see their family physician. We need to collect data from couples in order to assess the reliability of the information supplied by men regarding their partners pregnancy outcomes. We need to include questions that address the concerns of men regarding their mothers and their partners.

CONCLUSION

If the data and clinical impressions contained in this article and the deductions are correct, there appears to be a definable and diagnosable constellation of symptoms and problems comprising a syndrome, (Post Abortion Survivor Syndrome, PASS) that occurs in people whose parents have chosen to terminate other pregnancies. This constellation appears distinct from any combination of symptoms that might arise when a person survives the loss of a sibling by miscarriage. . Since many of the major symptoms arise from existential dilemmas, anyone attempting to diagnose this syndrome would need to ask appropriate questions.

Unless the conflicts behind these dilemmas are addressed it is unlikely that the expression of these conflicts, eg. difficult to treat depressions or the repeated need for psychiatric admission, will subside. This study needs to be replicated with a larger sample. In the meantime, it behoves the practising clinician to ask appropriate questions in an effort to determine whether a patient who presents with “depression” might be suffering from causes that are not biochemical or situational but rather from conflicts about life and living when a sibling was terminated.

References

1. Chodoff P. Late effects of the concentration camp syndrome. **Arch Gen Psych** 1963;8:323-333.
- 2, Niederland, WG. Clinical observations on the ‘Survivor Syndrome’. **Int J Psychoanalysis** 1968;49:313-15.
- 3, Poltawska W. **And I am Afraid of my Dreams**. New York: Hippocrene, 1989.
4. Kaminer H, Lavie P. Sleep and dreams in well adjusted and less adjusted Holocaust survivors. In: Stroebe MD, Stroebe W, O’Harrison R, eds. **Handbook of Bereavement**. New York: Cambridge University Press, 1994:331-345.
9. Rowland-Klein D, Dunlop R. The transmission of trauma across generations: identification with parental trauma in children of Holocaust survivors. **Aus NZ J Psych** 1997;31:358-369.
7. Krell R. Holocaust families: the survivors and their children. **Compr Psych** 1979;20:560-568.
5. . Barocas H, Barocas C. Manifestations of concentration camp effects on the second generation. **Am J Psych** 1973;103:820-821
8. Kestenberg JS. Survivor-parents and their children. In: Bergmann MS, Jucovy ME, eds. **Generations of the Holocaust**. New York: Columbia University Press, 1990:83-102.
6. . Danieli Y. Families of survivors of the Nazi Holocaust. Some short- and long-term effects. In: Spielberg CD et al., ed. **Stress and Anxiety**. Vol. 8. New York: McGraw Hill, 1981;405-421.
10. Miller A. **The drama of being a child**. 2nd ed. London: Virago, 1995.

11. Wardi D. **Memorial candles: children of the Holocaust.** (Goldblum N, trans.) London: Tavistock/Routledge, 1992.
 12. Ney PG. Transgenerational child abuse, **Child Psychiatry Hum Dev** 1988; 18:151-168.
 13. Ney PG. Child Mistreatment: Possible Reasons for its Transgenerational Transmission **Can J Psychiatry** 1989;34:594-601.
 14. Robinson S, Hemmendinger J, Netanel R, et al. Retraumatization of Holocaust survivors during the Gulf War and SCUD missile attacks on Israel. **Brit J Med Psych** 1994;67:353-362.
 15. Baider L, Peretz T, Hadani PE, et al. Transmission of response to trauma? Second generation Holocaust survivors' reaction to cancer. **Am J Psych** 2000;157(6):904-910.
 16. Wenzel T, Griengl H, Stompe T, et al. Psychological disorders in survivors of torture: exhaustion, impairment and depression. **Psychopathology** 2000;33:292-296.
 17. Bramsen I, van der Ploeg HM. Fifty years later: the long-term psychological adjustment of ageing World War II survivors. **Acta Psychiatr Scand** 1999;100:350-358.
 18. North CS, Nixon SJ, Shariat S, et al. Psychiatric disorders among survivors of the Oklahoma City bombing. **JAMA** 1999;282(8): 755-762.
 19. Bolton D, O’Ryan D, Udwin O, et al. The long-term psychological effects of a disaster experienced in adolescence: II: general psychopathology. **J Child Psychol, Psychiatr** 2000;41(4):513-523.
 20. Ney PG, Fung T, Wickett AR, et al. Effects of pregnancy loss on women’s health. **Soc Sci Med** 1994;38(9): 1193-2000.
 21. Ney PG, McPhee J, Moore C, et al. Child Abuse: A Study of the Child's Perspective. **Child Abuse Negl** 1986;10:511-518.
 22. Ney PG, Fung T, Wickett AR. Child Neglect: The Precursor to Child Abuse. **Pre- and Perinatal Psychology J.** 1993;8(2): 95-112.
 23. Kent I, Greenwood RD, Nicholls W. Emotional sequelae of elective abortion. **BC Med J** 1978;20:118-119
 24. Anonymous Patient Quotation. In: Ney PG, Peeters MA. **Abortion Survivors.** Victoria: Pioneer Publishing, 1996.
-

25. Sago H, Hayashi S, Saito M., The outcome and prognostic factors in twin-twin transfusion syndrome following fetoscopic laser surgery. **Prenat Diagn.** 2010 Dec; 30:1185-91.

-
26. Segal NL Suicidal behaviours in surviving monozygotic and dizygotic co-twins: is the nature of the co-twin's cause of death a factor? *Suicide Life Threat Behav.* 2009; 39: 569-75.
27. Rinborg, A, Lidgaard C, Freieslaeben NC et al Vanishing twins: a predictor of small-for-gestation age in IVF singletons. *Hum Reprod.* 2007 Oct; 22 (10) : 2707-14. Epub 2007 Aug:29.
28. Hayton A. www.wombtwin.com
29. . Ney PG, Fung T, Wickett AR, the Relationships between Induced Abortion and Child Abuse and Neglect: Four Studies, *Pre and Perinatal Psychology J.* 8: 43- 63 1993.
30. Ney PG, Ball K, Sheils C. Results of group psychotherapy for abuse, neglect and pregnancy loss. *Current Women's Health Review.* 2010; 6: 332-340.
- . 31. Kersting A, Ohrmann P, Neural Activation Underlying Acute Grief in Women After the Loss of an Unborn Child. *Am J. Psychiatry* 2009, 166: (12) 1402-1410
32. . Ney PG, Fung T, Wickett AR, et al. Effects of pregnancy loss on women's health. **Soc Sci Med** 1994;38(9): 1193-2000.

Table 1

How I found out about my mother's pregnancy loss

| | Frequency | Percent | Cumulative Percent |
|---------------------------|-----------|---------|-----------------------|
| no resp, N/A | 120 | 41.4 | 41.4 |
| mother told me | 97 | 33.4 | 74.8 |
| father told me | 5 | 1.7 | 76.6 |
| sibling told me | 7 | 2.4 | 79.0 |
| I asked | 12 | 4.1 | 83.1 |
| still not sure | 30 | 8.0 | 90.9 |
| fam. or friend told me | 4 | 1.4 | 92.1 |
| other way | 7 | 2.4 | 94.5 |
| discovered evidence | 5 | 1.7 | 96.2 |
| overheard conver. | 11 | 3.7 | 100.0 |
| Total | 290 | 100.0 | |

Table 2

How my children found out about my pregnancy loss

| | Freq | Percent | Cumulative % |
|-----------------------------|-------------|----------------|---------------------|
| I told them | 84 | 29 | 29 |
| My spouse told them | 17 | 5.9 | 34.9 |
| Sibling told them | 3 | 1 | 35.9 |
| My parent told them | 1 | 0.3 | 36.2 |
| A friend told them | 1 | 0.3 | 36.5 |
| A family member told them | 2 | 0.7 | 39.2 |
| They asked | 6 | 2.1 | 41.3 |
| They overheard conversation | 5 | 1.7 | 43 |
| They discovered evidence | 2 | 0.7 | 45.7 |
| They don't know | 40 | 13.8 | 69.9 |
| No response | 85 | 29.3 | 100 |

Table 3

Subjects' Present Problems and Symptoms

| | Standardized Coefficient | T | Sig |
|--|--------------------------|-------|------|
| Suicidal thoughts | | | |
| a) I am not glad to be alive | .357 | 5.552 | .000 |
| b) I feel life is not worth living | .259 | 4.355 | .000 |
| c) I hear or see or feel things that appear from nowhere | .213 | 4.269 | .000 |
| Repeated depression | | | |
| a) I am bothered by thoughts I can't control | .282 | 4.742 | .000 |
| b) I feel sad | .218 | 3.312 | .001 |
| c) My human relationships are poor | .167 | 2.561 | .011 |

Table 4

Symptoms most closely associated with “I feel I don’t deserve to be alive” (PASS)

| | Untandardized Coefficients B. | T | Sig |
|---|-------------------------------|-------|------|
| a) I am not glad to be alive | .428 | 7.117 | .000 |
| b) I feel something terrible is going to happen to me | .202 | 4.772 | .000 |
| c) I have injured myself | .141 | 3.521 | .000 |
| d) I fear I am losing my mind | .172 | 3.481 | .001 |
| e) I have tried to kill myself | .128 | 2.403 | .017 |

Table 5

Existential guilt and all pregnancy outcomes of subjects' mother's first pregnancy

Included Variables

| <u>Dependant variable:</u> I feel I don't deserve to be alive | Unstandardized Coefficient, B | t | Sig |
|--|----------------------------------|-------|------|
| Total number of abortions in mothers' first pregnancy | .013 | 2.254 | .025 |

Excluded Variables:

| | Beta In | t | Sig. |
|--|---------|--------|------|
| Total number of full term, normal weight in first pregnancy of mother | -.042 | -.673 | .501 |
| Total number of full term, low birth weight in first pregnancy of mother | .065 | 1.111 | .268 |
| Total number of premature in first pregnancy of mother | .074 | 1.257 | .210 |
| Total number of miscarriage in first pregnancy of mother | -.051 | -.875 | .383 |
| Total number of stillbirth in first pregnancy of mother | -.018 | -.314 | .754 |
| Early infant death in first pregnancy of mother | -.061 | -1.040 | .299 |

Step wise regression with cut off $p < 0.05$

Table 6

Partial correlation coefficients of 16 Symptoms

| | Glad | Know | Pleased | Ability | Relation | Worth | Injured | Kill | Terrible | Deserve | Sad | Angry | Unreal | Mind | Hear |
|----------|--------|--------|---------|---------|----------|--------|---------|--------|----------|---------|--------|--------|--------|--------|--------|
| Know | .583** | | | | | | | | | | | | | | |
| Pleased | .612** | .764** | | | | | | | | | | | | | |
| Ability | .447** | .562** | .646** | | | | | | | | | | | | |
| Relation | .525** | .565** | .620** | .555** | | | | | | | | | | | |
| Worth | .569** | .414** | .482** | .301** | .404** | | | | | | | | | | |
| Injured | .246** | .199* | .224** | .207** | .217** | .220** | | | | | | | | | |
| Kill | .121 | .111 | .126 | .111 | .177* | .185* | .470** | | | | | | | | |
| Terrible | .308** | .338** | .439** | .400** | .389** | .364** | .275** | .238** | | | | | | | |
| Deserve | .575** | .382** | .467** | .431** | .456** | .415** | .392** | .331** | .485** | | | | | | |
| Sad | .549** | .490** | .566** | .467** | .599** | .423** | .285** | .189* | .459** | .454** | | | | | |
| Angry | .501** | .517** | .541** | .460** | .535** | .361** | .256** | .206** | .346** | .424** | .688** | | | | |
| Unreal | .322** | .309** | .356** | .364** | .418** | .293** | .382** | .233** | .464** | .422** | .453** | .462** | | | |
| Mind | .538** | .464** | .570** | .421** | .501** | .414** | .255** | .238** | .435** | .538** | .554** | .550** | .585** | | |
| Hear | .169** | .104 | .098 | .143 | .148 | .098 | .248** | .181* | .227** | .225** | .162* | .168** | .371** | .325** | |
| Control | .416** | .449** | .471** | .431** | .474** | .354** | .233** | .201* | .424** | .400** | .490** | .466** | .521** | .617** | .356** |

* p < 0.01 ** P< 0.001 Controlling for age, sex, marital status and number of children.

Glad = I am **not glad** to be alive, Know = I **don't know** who I am, I am **not pleased** with who I am, I am not using my **abilities**, My human **relationships** are insecure, I feel life is not **worth** living, I have **injured** myself, I have tried to **kill** myself, I sense something **terrible** is going to happen to me, I feel I don't **deserve** to be alive, I feel **sad**, I feel **angry**, I have feelings that things are **unreal**, I fear I am losing my **mind**, I **hear** or see or feel things that appear from nowhere. I am bothered by thoughts I can't **control**

TABLE 7
Most prominent symptoms associated with abortions of the mothers of the subjects.

| Dependant variable in response to the following questions | Unstandardized Coefficient, B | t | Sig. |
|--|-------------------------------|--------|------|
| 1. Did your parents consider aborting you? Yes, Maybe, No | | | |
| a) I have feelings that things are unreal | .065 | 2.377 | .018 |
| b) I have tried to kill myself | .076 | 2.183 | .030 |
| 2. Poor chances of survival before I was born (0 - 8) | | | |
| a) I have feelings that things are unreal | .355 | 3.210 | .001 |
| b) I am not pleased with who I am | .213 | 2.078 | .039 |
| 3. Very upset when found out about pregnancy loss (0 - 8) | | | |
| a) I know who I am | -.251 | -2.935 | .004 |
| b) I have feelings things are unreal | .202 | 2.301 | .022 |
| c) I feel I don't deserve to be alive | .182 | 2.076 | .039 |
| 4. Very upset about mother's pregnancy loss now (0 - 8) | | | |
| a) I feel I don't deserve to be alive | .214 | 2.715 | .007 |
| 5. Statistically my chance of being aborted were very high (0-8) | | | |
| a) I have tried to kill myself | .337 | 5.414 | .000 |
| 6. Total # of abortions in preg. 1-9 of subject's mother | | | |
| a) I am bothered by thoughts I can't contro | 145 | 2.486 | .013 |

Step wise regression; cut off set at $p < .05$

There were no significant associations of symptoms for premature births, still births or early infant deaths.

Table 8
Past Problems and Abortions of the Mothers of the Subjects

| <u>Dependant variables are:</u> | Untandardized coefficient B | T | Sig |
|--|-----------------------------|-------|------|
| 1. # abortions in first pregnancy | | | |
| a) Psychiatric hospitalization | .095 | 2.012 | .045 |
| 2. # person's whose mother had at least one abortion | | | |
| a) Difficulty being a good parent | .136 | 3.037 | .003 |
| b) Suicidal thoughts | .123 | 2.798 | .005 |
| 3. # abortions in pregnancy 1-9 of subject's mother | | | |
| a) Suicidal thoughts | .186 | 2.517 | .012 |
| b) Difficulty being a good parent | .188 | 2.505 | .013 |
| 4. Feel poorly about mother's pregnancy loss now | | | |
| a) Poor physical health | .971 | 2.553 | .011 |
| 5. High statistical chance of being aborted. | | | |
| a) Substance abuse | .688 | 2.885 | .004 |
| b) Psychiatric hospitalization | .931 | 2.292 | .023 |
| c) Suicidal thoughts | .463 | 2.012 | .045 |

Step wise regression with cut off at $p. < .05$

There were no significant problems in the subject's past associated with miscarriages in the subject's mother's first pregnancy. There were no significantly associated past problems for those whose mother never had an abortion but had some other pregnancy loss.

Table 9
Transgenerational aspects of abortion, subject and her mother.

Controlling for: Age, Sex, Marital Status

| | M_Abo_1 1. | M_Abo_2 2. | M_Abo_T 3. | T_M_Abo 4. | My_Abo_1 5. | My_Abo_ 6. |
|----------|---------------|---------------|---------------|---------------|----------------|---------------|
| M_Abo_2 | .157* | | | | | |
| M_Abo_T | .474** | .450** | | | | |
| T_M_Abo | .514* | .381** | .820** | | | |
| My_Abo_1 | .174* | -.054 | .142* | .193* | | |
| My_Abo_2 | .028 | .013 | .108 | .104 | .286** | |
| T_My_Abo | .110 | .059 | .147 | .173* | .777** | .478** |

* p.<.05 **p<.01 Partial correlation coefficients controlling for age, sex and marital status.

- 1) Total number of abortions in the first pregnancy of subjects' mothers (M_Abo_1)
- 2) Total number of abortions in the second pregnancy of subjects' mothers (M_Abo_2)
- 3) Total numbers of abortions in any pregnancy for all subject's mothers (T_M_Abo)
- 4) Total numbers of mothers of subjects who had one or more abortions (M_Abo_T)
- 5) Total number of abortions in the first pregnancy of the subject (My_Abo_1)
- 6) Total number of abortions in the second pregnancy of the subject (My_Abo_2)
- 7) Total number of subjects who had one or more abortions (T_My_Abo)

Table 10

Most prominent symptoms in descending order of impact for PASS.

| | |
|--|--------------------------|
| 1. I feel I don't deserve to be alive | Existential guilt |
| 2. I am not glad to be alive | Existential sorrow |
| 3. Something terrible is going to happen to me | Sense of impending doom |
| 4. I have tried to kill myself | Self destructive |
| 5. I fear I am losing my mind | Tenuous grasp of reality |
| 6. I have injured myself | Self injurious |
| 7. I feel sad | Sorrow |
| 8. I am not pleased with who I am | Low self esteem |
| 9. I have feelings that things are unreal | Dissociation |
| 10. I am bothered by thoughts I cannot control | Obsessive thinking |
| 11. I do not know who I am | Poor self identity |
| 12. I feel life is not worth living | Depression |

9 - 12 from a step wise regression of answers to questions regarding statistical chance of being aborted.
